

Polskie Towarzystwo Statystyczne
Oddział we Wrocławiu

ŚLĄSKI PRZEGLĄD STATYSTYCZNY

Silesian Statistical Review

Nr 14 (20)



Wydawnictwo Uniwersytetu Ekonomicznego we Wrocławiu
Wrocław 2016

RECENZENCI WSPÓŁPRACUJĄCY Z CZASOPISMEM

Milan Bašta, Tadeusz Borys, Mariusz Czekala, Jakub Fisher, Ewa Frątczak, Stanisława Hronová, Helena Jasiulewicz, Alina Jędrzejczak, Wojciech Kordecki, Ryszard Kryszewski, Dorota Kuchta, Jitka Langhamrová, Tomáš Loster, Ivana Malá, Krystyna Melich, Zofia Mielecka-Kubień, Witold Miszczak, Juliusz Siedlecki, Jaroslav Sixta, Włodzimierz Szkutnik, Jerzy Wawrzynek, Witold Więśław, Jiří Witzany, Emília Zimková

RADA NAUKOWA

Walenty Ostasiewicz (Uniwersytet Ekonomiczny we Wrocławiu, Polska)

Tadeusz Bednarski (Uniwersytet Wrocławski, Polska)

Ivan Belko (Belarusian State University, Belarus)

Luisa Canal (University of Trento, Italy)

Karlheinz Fleischer (Philipps-Universität Marburg, Germany)

Francesca Greselin (University of Milano-Bicocca, Italy)

Stanisław Heilpern (Uniwersytet Ekonomiczny we Wrocławiu, Polska)

Stanislava Hronová (VSE Prague, the Czech Republic)

Salvatore Ingrassia (University of Catania, Italy)

Jerzy Śleszyński (Uniwersytet Warszawski, Polska)

Halina Woźniak (Urząd Statystyczny we Wrocławiu, Polska)

Michele Zenga (University of Milano-Bicocca, Italy)

Emília Zimková (Matej Bel University Banská Bystrica, Slovakia)

Ricardas Zitikis (University of Western Ontario, Canada)

KOMITET REDAKCYJNY

Zofia Rusnak (Uniwersytet Ekonomiczny we Wrocławiu, Polska) –
redaktor naczelny

Katarzyna Ostasiewicz (Uniwersytet Ekonomiczny we Wrocławiu, Polska)

Angiola Pollastri (University of Milano-Bicocca, Italy)

Grażyna Trzpiot (Uniwersytet Ekonomiczny w Katowicach, Polska)

Reinhard Viertl (Vienna University of Technology, Austria)

Edyta Mazurek – sekretarz

edyta.mazurek@ue.wroc.pl

+48 71 71 36 80 325

Spis treści

Aims and scope 5

- Oscar Sheynin:** On the history of university statistics 7
- Marian Matloka:** h -Preinvex fuzzy processes 27
- Joanna Dębicka, Beata Zmyślona:** Construction of multi-state life tables for critical illness insurance – influence of age and sex on the incidence of health inequalities 41
- Wiktor Ejsmont:** Podstawowe pojęcia wolnej probabilistyki 65
- Edyta Mazurek:** Podatek dochodowy w kontekście rodziny 75
- Katarzyna Ostasiewicz:** Kto co konsumuje i czy wystarczająco dużo: gospodarka i bieda, czyli Nagroda imienia Nobla z dziedziny ekonomii dla Angusa Deatona (2015) 89
- Agnieszka Thier:** Analiza sposobów pomiaru oraz skutków deficytu zasobów wodnych na świecie 111
- Damian Gąska:** Wykorzystanie sieci bayesowskich do prognozowania bankructwa firm 131
- Walenty Ostasiewicz:** Metabometria 145
- Monika Hadaś-Dyduch:** Iluzja, marzenia a rzeczywistość – bezpośrednia i niebezpośrednia inwestycja w indeksy giełdowe na przykładzie produktów inwestycyjnych 185
- Agnieszka Marciniuk:** 23. Scientific Statistical Seminar “Wrocław-Marburg” 203
- 23. Scientific Statistical Seminar “Wrocław-Marburg”, Pottenstein-Kirchenbirkig, 28.09.2015 – 1.10.2015.**
Extended Abstracts 207
- Beata Zmyślona:** Application of Mathematics and Statistics in Economics. The 18th International Scientific Conference 229
- Tadeusz Gerstenkorn:** Włodzimierz Kryszicki matematyk-stochastyk (1905–2001) 233
- Walenty Ostasiewicz:** Profesor Ryszard Antoniewicz (19.08.1939 – 20.02.2015) 243
- Walenty Ostasiewicz:** Nobel, Non Nobel, Ig Nobel, and Alternative Nobel Prizes 251
- Agata Girul:** Ważniejsze dane społeczno-gospodarcze o województwach 255

Summaries

- Oscar Sheynin:** On the history of university statistics 7
- Marian Matłoka:** h-Preinvex fuzzy processes 27
- Joanna Dębicka, Beata Zmyślona:** Construction of multi-state life tables for critical illness insurance – influence of age and sex on the incidence of health inequalities 41
- Wiktor Ejsmont:** Basic concepts of free probability theory 73
- Edyta Mazurek:** The income tax in the context of the family 87
- Katarzyna Ostasiewicz:** Who consumes what and is it enough: economy and poverty. Prize in Economic Sciences in Memory of Alfred Nobel for Angus Deaton (2015) 110
- Agnieszka Thier:** Analysis of ways of measurement and the consequences of water shortage in the world 128
- Damian Gąska:** Bankruptcy prediction with Bayesian networks 143
- Walenty Ostasiewicz:** Metabometrics 182
- Monika Hadaś-Dyduch:** The illusion, dreams and reality – direct and indirect investment in stock indices on the example of investment products 201

Aims and scope

Aims and scope of this journal were determined already in the period of the historical changes that took place in 1989 in the Europe, which had a great meaning for Poland, especially for the subsequent political and economic transformations. The introduction of the democratic system, and the transition from the state-controlled economy to the free market one were the driving forces behind the new Polish economy.

In the early 1990s, Poland made great progress towards achieving a fully democratic government and a market economy. In November 1990, Lech Wałęsa was elected President for a 5-year term. In 1991 were held the first free parliamentary elections. In the same year, 1991, the first issue of the journal was published under the title *Statistical Review of Lower and Opole Silesia*. In the foreword of that first issue it was stated what follows. “The changes in the socio-economic life of Lower Silesia and Opole region caused the Council of Wrocław Branch of Polish Statistical Society to publish Statistical Review of Lower and Opole Silesia, starting from the year 1991. This idea could come to life thanks to the generous help of directors of Voivodeship Statistical Offices in Jelenia Góra, Legnica, Wałbrzych and Wrocław, with a special involvement of the director of Statistical Office in Wrocław”. The initial goal of the founders of the journal was to dedicate the journal to “ecological problems, demographic issues as well as social and economic well-being”.

Starting in the year 2002 the journal has been published with a new layout and under a new title: *Silesian Statistical Review*. Together with *Statistical Review (Przegląd Statystyczny)* and *Statistical News (Wiadomości Statystyczne)*, *Silesian Statistical Review* is now one of the three major journals in Poland dedicated to general statistical problems. Special attention has been focused on general methodological issues, as well as on the applications of various statistical methods in solving real social and economic problems. Papers concerning all topics of quality of life are published regularly. Historical essays are included on regular basis.

After 25 years of the existence, by entering in the next quarter of the century of its existence with the issue of 2016, the main scope of journal is amplified. This is again caused by changes which took place on the

whole planet. In order to meet the challenge mounted by dramatic consequences of human dominance over the planet the scope of journal has been amplified to include any problems concerning the quality of human life, respecting all other forms of lives and not compromising the possibilities for future generations to live their ways of life.

Starting from the year 2016, *Silesian Statistical Review* is considered as a *Journal of Oikometrics*

The name, derived from Greek words *οικος* and *μετρο*, suggests that the journal focus is upon Nature's house (*oikos*), as a subject matter of a study, and the measurement, as a prevailing methodology of study. The journal is treated as an *interdisciplinary forum on a sustainable livelihood*. Contrary to the inscription on the door of Plato's Academy: *let no one ignorant of geometry enter here*, over the door to *Journal of Oikometrics* there is hanged the signboard with the inscription: *Everyone who cares about, and interested in any issue of sustainable livelihood is welcomed here*.

The Journal welcomes therefore papers from specialists in sustainability science, ecology, ecological economics and any other alternatives to neoclassical economics. It encompasses – but is not limited to – the following topics:

- actuarial methods and their applications,
- social justice, inequality, polarization, and stratification,
- quality of institutional performance,
- social metabolism, its measurement and analysis,
- statistical education,
- sustainable development,
- environmentalism.

As the official journal of the Polish Statistical Society, Branch in Wrocław, it is designed also to attract papers that have direct relation with the activity of the Society, particularly in the field of education, promotion and rising awareness of the statistics role in the civilization development.

Walenty Ostasiewicz

Marian Matłoka

Poznań University of Economics and Business

ISSN 1644-6739
e-ISSN 2449-9765

DOI: 10.15611/sps.2016.14.02

Summary: We introduce the notion of *h*-preinvex fuzzy processes. We study their properties and give some inequalities of Hadamard-type for *h*-convex fuzzy processes.

Keywords: *h*-preinvex fuzzy process, Hadamard's inequalities.

1. Introduction

Multifunctions have many diverse and interesting applications in control problems and the theory of contingent equations, in mathematical economics, and in various branches of analysis (for example, see [Aumann 1965; Brunovsky 1968; Debreu 1983; Hermes 1968]).

The special class of the multifunctions is called convex processes. Convex processes were studied first by R.T. Rockafellar who was interested in extending properties of linear transformations to a large class of maps preserving convexity and which arise naturally in economic theory. He elaborated a duality theory for these convex processes and developed his study in his book [Rockafellar 1970]. P. Coutat studied continuous closed convex sets and gave many new properties of convex processes [Coutat 1996].

The extension of this notion to the fuzzy framework was done by M. Matłoka [2000]. Y.R. Syau, C.Y. Low and T.H. Wu [2002] observed that the Matłoka definition is very strict. Therefore, they give another definition that extends the Matłoka definition. In recent years, many generalizations of convexity have appeared in the literature aiming at applications to duality theory and optimality conditions. In 1997 R. Pini and Ch. Singh [20] introduced (Φ_1, Φ_2) -convex functions and studied some of their properties [Pini, Singh 1997]. They showed that some of the well-known classes of generalized convex functions (e.g. B-vex functions [Bector, Singh 1991], invex functions [Hanson 1981]) form subclasses of the class of (Φ_1, Φ_2) -convex functions. In 1999 E.A. Youness showed that many results for convex functions actually hold for a wider class of functions, called E-convex functions [Youness 1999]. In 1978,

W.W. Breckner introduced s -convex functions as a generalization of convex functions [Breckner 1978] and in 1993 studied the set valued version [Breckner 1993]. The extension of this notions to the fuzzy framework was done by M. Matłoka [1999, 2012], and Y. Chalco-Cano and M.A. Rojas-Medar [2004]. In 2007 S. Varošanec [26] introduced a large class of functions, the so-called h -convex functions [Varošanec 2007]. This class contains several well-known classes of functions such as convex functions and s -convex functions.

In this work, we introduce a new class of fuzzy processes, the so-called h -preinvex fuzzy processes. This class contains several well-known classes of fuzzy processes such as convex and s -convex.

The plan of the paper is as follows. In Section 2, we introduce the notations and definitions used throughout the paper. In Section 3, we establish the main results and finally, in Section 4, we show some, properties and the connection with fuzzy integral for fuzzy set-valued maps.

2. Preliminaries

Let R^n denote the n -dimensional Euclidean space.

A fuzzy subset of R^n is a function $u: R^n \rightarrow [0, 1]$. Let $\mathcal{F}(R^n)$ denote the set of all nonempty fuzzy sets in R^n . A fuzzy set u is called convex [Lowen 1980], if

$$u(ty_1 + (1 - t)y_2) \geq \min\{u(y_1), u(y_2)\},$$

for all $y_1, y_2 \in \text{supp}(u) = \overline{\{y: u(y) > 0\}}$ and $t \in (0, 1)$.

We shall define addition and scalar multiplication on $\mathcal{F}(R^n)$ by the usual extension principle, i.e. for $u, v \in \mathcal{F}(R^n)$ and $\lambda \in R$, $u + v$ and $\lambda \cdot u$ are defined for any $y \in R^n$ by

$$(u + v)(y) = \sup_{y_1, y_2: y_1 + y_2 = y} \min(u(y_1), v(y_2))$$

and

$$(\lambda u)(y) = \begin{cases} u\left(\frac{y}{\lambda}\right) & \text{if } \lambda \neq 0, \\ \mathcal{X}_{\{0\}}(y) & \text{if } \lambda = 0, \end{cases}$$

where for any subset $A \subseteq R^n$, \mathcal{X}_A denotes the characteristic function of A .

We can define an order \subseteq on $\mathcal{F}(R^n)$ by setting

$$u \subseteq v \Leftrightarrow u(y) \leq v(y), \forall y \in R^n.$$

Let $u \in \mathcal{F}(R^n)$. For $0 < \alpha \leq 1$, we denote $[u]^\alpha = \{y \in R^n: u(y) \geq \alpha\}$ the α -level set of u . $[u]^0 = \text{supp}(u) = \{y: u(y) > 0\}$ is called the support of u .

A fuzzy set $u: R^n \rightarrow [0, 1]$ is said to be a fuzzy compact set if $[u]^\alpha$ is compact for all $\alpha \in [0, 1]$. For any fuzzy compact sets u and v it is verified that

- $u \subseteq v \Leftrightarrow [u]^\alpha \subseteq [v]^\alpha, \forall \alpha \in [0, 1]$,
- $[\lambda u]^\alpha = \lambda [u]^\alpha, \forall \lambda$ and $\forall \alpha \in [0, 1]$,
- $[u + v]^\alpha = [u]^\alpha + [v]^\alpha, \forall \alpha \in [0, 1]$.

Definition 2.1. A fuzzy set $u: R^n \rightarrow [0, 1]$ is called *h-convex fuzzy set* iff

$$u(h(t)x_1 + h(1 - t)x_2) \geq \min\{u(x_1), u(x_2)\}$$

for all $x_1, x_2 \in R^n, t \in (0, 1)$ where $h: J \rightarrow R$ is a non-negative function, $h \not\equiv 0$ and J is an interval, $(0, 1) \subset J$.

Definition 2.2. [Weir, Mond 1988] Let X be a nonempty set in R^n and $x \in X$.

Then the set X is said to be *invex at x* with respect to $\eta: X \times X \rightarrow R^n$, if

$$x + t\eta(y, x) \in X$$

for all $y \in X$ and $t \in [0, 1]$.

X is said to be an *invex set* with respect to η , if X is invex at each $x \in X$.

Definition 2.3. A fuzzy set $u: X \times R^n \rightarrow [0, 1]$ is called *(η, h)-convex fuzzy set* iff

$$u(x_1 + t\eta(x_2, x_1), h(t)y_1 + h(1 - t)y_2) \geq \min\{u(x_1, y_1), u(x_2, y_2)\}$$

for all $x_1, x_2 \in X, y_1, y_2 \in R^n$ and $t \in [0, 1]$.

Definition 2.4. The graph of a mapping $F: X \rightarrow \mathcal{F}(R^n)$, denoted by G_F , is a fuzzy set in $X \times R^n$ such that for any $(x, y) \in X \times R^n$

$$G_F(x, y) = F(x)(y).$$

Definition 2.5. The function f on the invex set X is said to be *preinvex with respect to η* , if

$$f(x + t\eta(y, x)) \leq h(t)f(x) + h(1 - t)f(y)$$

for each $x, y \in X$ and $t \in (0, 1)$.

Let us note that:

- if $\eta(y, x) = y - x$ then we get the definition of h -convex function introduced by S. Varošanec [2007],
- if $h(t) = t$ then our definition reduces to the definition of preinvex function [Weir, Mond 1988],
- if $\eta(y, x) = y - x$ and $h(t) = t$ then we obtain the definition of convex function,
- if $\eta(y, x) = y - x$ and $h(t) = t^s$ then our definition reduces to the definition of s -convex function [Breckner 1978].

3. h -Preinvex fuzzy processes

In this section, we present some basic properties of h -preinvex fuzzy processes.

Definition 3.1. A mapping F from invex set $X \subset R^m$ into $\mathcal{F}(R^n)$ is called h -preinvex fuzzy process iff for any $x_1, x_2 \in X$ and $t \in (0, 1)$ it satisfies the condition

$$h(t)F(x_1) + h(1-t)F(x_2) \subseteq F(x_1 + t\eta(x_2, x_1)),$$

where $h: J \rightarrow R$ is a positive function and J is an interval, $(0, 1) \subset J$.

If $\eta(x_2, x_1) = x_2 - x_1$ then such a mapping we will call h -convex fuzzy process.

Remark 3.1. If $\eta(x_2, x_1) = x_2 - x_1$ and $h(t) = t^s$ then our definition reduces to the definition of s -convex fuzzy process defined by Y. Chalco-Cano and M.A. Rojas-Medar [2004].

Example 3.1. Consider $F: X \rightarrow \mathcal{F}(R)$ defined by

$$F(x) = \chi_{[f(x), \infty]}$$

where $f(x)$ is h -preinvex function from X into R . Then F is h -preinvex process.

Theorem 3.1. A mapping $F: X \rightarrow \mathcal{F}(R^n)$ is a h -preinvex fuzzy process if and only if

$$\begin{aligned} & F(x_1 + t\eta(x_2, x_1))(y) \\ & \geq \sup_{y_1, y_2: h(t)y_1 + h(1-t)y_2 = y} \min \{F(x_1)(y_1), F(x_2)(y_2)\} \end{aligned}$$

for all $x_1, x_2 \in X$ and $y \in R^n$.

Proof. Suppose that F is *h*-preinvex fuzzy process. Let $x_1, x_2 \in X$, $t \in (0, 1)$ and $y \in R^n$. Then, from Definition 3.1 we have

$$\begin{aligned} F(x_1 + t\eta(x_2, x_1))(y) &\geq (h(t)F(x_1) + h(1-t)F(x_2))(y) \\ &= \sup_{y_1, y_2: y_1 + y_2 = y} \min \{h(t)F(x_1)(y_1), h(1-t)F(x_2)(y_2)\} \\ &= \sup_{y_1, y_2: y_1 + y_2 = y} \min \left\{ F(x_1)\left(\frac{y_1}{h(t)}\right), F(x_2)\left(\frac{y_2}{h(1-t)}\right) \right\} \\ &= \sup_{y_1, y_2: h(t)y_1 + h(1-t)y_2 = y} \min \{F(x_1)(y_1), F(x_2)(y_2)\}. \end{aligned}$$

Conversely, let us suppose that the inequality from the theorem is satisfied. Then for all $x_1, x_2 \in X$, $t \in (0, 1)$ and $y \in R^n$, we have

$$\begin{aligned} F(x_1 + t\eta(x_2, x_1))(y) &\geq \sup_{y_1, y_2: h(t)y_1 + h(1-t)y_2 = y} \min \{F(x_1)(y_1), F(x_2)(y_2)\} \\ &= \sup_{y'_1, y'_2: y'_1 + y'_2 = y} \min \left\{ F(x_1)\left(\frac{y'_1}{h(t)}\right), F(x_2)\left(\frac{y'_2}{h(1-t)}\right) \right\} \\ &= \sup_{y'_1, y'_2: y'_1 + y'_2 = y} \min \{h(t)F(x_1)(y'_1), h(1-t)F(x_2)(y'_2)\} \\ &= (h(t)F(x_1) + h(1-t)F(x_2))(y) \end{aligned}$$

which implies that F is *h*-preinvex fuzzy process.

Theorem 3.2. Let $F: R^m \rightarrow \mathcal{F}(R^n)$ be a mapping such that

- (1) $F(x_1 + x_2) \supseteq F(x_1) + F(x_2), \quad \forall x_1, x_2 \in R^m,$
- (2) $F(tx) = h(t)F(x), \quad \forall t > 0$ and $\forall x \in R^m.$

Then, F is *h*-preinvex fuzzy process.

Proof. Let $x_1, x_2 \in R^m$, $t \in (0, 1)$ and $y \in R^n$. Then, from the addition and scalar multiplication on $\mathcal{F}(R^n)$, and from the conditions (1) and (2), we have

$$\begin{aligned} F(tx_1 + (1-t)x_2)(y) &\geq (F(tx_1) + F((1-t)x_2))(y) \\ &= \sup_{y_1, y_2: y_1 + y_2 = y} \min \{F(tx_1)(y_1), F((1-t)x_2)(y_2)\} \\ &= \sup_{y'_1, y'_2: h(t)y'_1 + h(1-t)y'_2 = y} \min \{F(tx_1)(h(t)y'_1), F((1-t)x_2)(h(1-t)y'_2)\} \end{aligned}$$

$$\begin{aligned}
 &= \sup_{y'_1, y'_2: h(t)y'_1 + h(1-t)y'_2 = y} \min \{ (h(t)F(x_1))(h(t)y'_1), (h(1-t)F(x_2))(h(1-t)y'_2) \} \\
 &= \sup_{y'_1, y'_2: h(t)y'_1 + h(1-t)y'_2 = y} \min \{ F(x_1)(y'_1), F(x_2)(y'_2) \}
 \end{aligned}$$

which means that F satisfies the inequality from the Theorem 1. Therefore, F is h -preinvex fuzzy process.

Theorem 3.3. If $F: R^m \rightarrow \mathcal{F}(R^n)$ is a h -preinvex fuzzy process and for any $t \in (0, 1)$ and $x \in X$ $x = x + t\eta(x, x)$ then for any $x \in X$, $F(x)$ is a h -convex fuzzy set.

Proof. Let $x \in X$. Since $x = x + t\eta(x, x)$ for all $t \in (0, 1)$, by the definition of h -preinvex fuzzy processes, we have for all $y_1, y_2 \in R^n$ and $t \in (0, 1)$,

$$\begin{aligned}
 &F(x)(h(t)y_1 + h(1-t)y_2) \\
 &= F(x + t\eta(x, x))(h(t)y_1 + h(1-t)y_2) \\
 &\geq \sup_{\substack{y'_1, y'_2: \\ h(t)y'_1 + h(1-t)y'_2 = h(t)y_1 + h(1-t)y_2}} \min \{ F(x)(y'_1), F(x)(y'_2) \} \\
 &\geq \min \{ F(x)(y_1), F(x)(y_2) \}.
 \end{aligned}$$

Hence, $F(x)$ is a h -convex fuzzy set. This completes the proof.

Theorem 3.4. The graph of h -preinvex fuzzy process is a (η, h) -convex fuzzy set in $X \times R^n$.

Proof. Let $x_1, x_2 \in X$ and $y_1, y_2 \in R^n$ and $t \in (0, 1)$.

Then we have

$$\begin{aligned}
 &G_F(x_1 + t\eta(x_2, x_1), h(t)y_1 + h(1-t)y_2) \\
 &= F(x_1 + t\eta(x_2, x_1))(h(t)y_1 + h(1-t)y_2) \\
 &\geq \sup_{\substack{y'_1, y'_2: \\ h(t)y'_1 + h(1-t)y'_2 = h(t)y_1 + h(1-t)y_2}} \min \{ F(x_1)(y'_1), F(x_2)(y'_2) \} \\
 &\geq \min \{ F(x_1)(y_1), F(x_2)(y_2) \} = \min \{ G_F(x_1, y_1), G_F(x_2, y_2) \}.
 \end{aligned}$$

So, G_F is a (η, h) -convex fuzzy set in $X \times R^n$.

Theorem 3.5. If F is a h -preinvex fuzzy process then for any $\alpha \in (0, 1]$

$$[F(x_1 + t\eta(x_2, x_1))]^\alpha \supset h(t)[F(x_1)]^\alpha + h(1-t)[F(x_2)]^\alpha$$

for any $x_1, x_2 \in X$ and $t \in (0, 1)$.

Proof. Let $x_1, x_2 \in X$ and $t \in (0, 1)$, and $\alpha \in (0, 1]$. Then we have

$$\begin{aligned}
 [F(x_1 + t\eta(x_2, x_1))]^\alpha &= \{y \in R^n: F(x_1 + t\eta(x_2, x_1))(y) \geq \alpha\}, \\
 [F(x_1)]^\alpha &= \{y_1 \in R^n: F(x_1)(y_1) \geq \alpha\}, \\
 [F(x_2)]^\alpha &= \{y_2 \in R^n: F(x_2)(y_2) \geq \alpha\}.
 \end{aligned}$$

Moreover

$$\begin{aligned}
 h(t)[F(x_1)]^\alpha + h(1-t)[F(x_2)]^\alpha &= \{y = h(t)y_1 + h(1-t)y_2: \\
 & y_1 \in [F(x_1)]^\alpha, y_2 \in [F(x_2)]^\alpha\}
 \end{aligned}$$

and

$$F(x_1 + t\eta(x_2, x_1))(y) \geq \sup_{\substack{y_1, y_2: \\ h(t)y_1 + h(1-t)y_2 = y}} \min \{F(x_1)(y_1), F(x_2)(y_2)\}.$$

This means that if $y \in h(t)[F(x_1)]^\alpha + h(1-t)[F(x_2)]^\alpha$ then $y \in [F(x_1 + t\eta(x_2, x_1))]^\alpha$, i.e.

$$[F(x_1 + t\eta(x_2, x_1))]^\alpha \supset h(t)[F(x_1)]^\alpha + h(1-t)[F(x_2)]^\alpha$$

what completes the proof.

4. Hadamard’s inequality

In this section, we present some inequalities of Hadamard type for *h*-preinvex fuzzy processes.

Let $\mathcal{F}_K(R^n)$ ($\mathcal{F}_C(R^n)$) denote the set of all fuzzy compact (compact and convex) sets and $K(R^n)$ the set of all nonempty, compact (crisp) subsets of R^n .

R.L. Aumann [1965] introduced the following definition of the integral of set-valued function.

Definition 4.1. The integral of a measurable set-valued function $G: [a, b] \rightarrow K(R^n)$ is defined by

$$\int_a^b G dt = \left\{ \int_a^b g(t) dt : g \in S(G) \right\},$$

where $\int_a^b g(t) dt$ is the Bochner-integral and $S(G)$ is the set of all integrable selectors of G , i.e.,

$$S(G) = \{g \in L^1([a, b], R^n): g(t) \in G(t) \text{ a. e.}\}.$$

For a mapping $F : [a, b] \rightarrow \mathcal{F}_K(R^n)$ let us define a mapping $F_\alpha : [a, b] \rightarrow K(R^n)$ by $F_\alpha(x) = [F(x)]^\alpha, \forall \alpha \in [0, 1]$.

Definition 4.2 [Puri, Ralescu 1987]. *A mapping F is called measurable (integrably bounded) if F_α is measurable (integrably bounded) for all $\alpha \in [0, 1]$.*

Proposition 4.1 [Puri, Ralescu 1987]. If $F : [a, b] \rightarrow \mathcal{F}_K(R^n)$ is integrably bounded, then there exists a unique fuzzy set $u \in \mathcal{F}_K(R^n)$ such that $[u]^\alpha = \int_a^b F_\alpha dt, \forall \alpha \in [0, 1]$.

The fuzzy set $u \in \mathcal{F}_K(R^n)$ in the above Proposition defines integral of F by $\int_a^b F dt = u$ iff $[u]^\alpha = \int_a^b F_\alpha dt$, for every $\alpha \in [0, 1]$.

Many important inequalities have been established for the class of convex functions, but the most famous is the Hermite–Hadamard’s inequality. These double inequalities are stated as:

(a) for convex function

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2},$$

(b) for s – convex function (see [Dragomir, Fitzpatrick 1999])

$$2^{s-1} f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{s+1},$$

(c) for h -convex function (see [Sarikaya et al. 2008])

$$\frac{1}{2h\left(\frac{1}{2}\right)} f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \int_0^1 h(t) dt [f(a) + f(b)],$$

(d) for preinvex function (see [Noor 2009])

$$f\left(a + \frac{1}{2}\eta(b, a)\right) \leq \frac{1}{\eta(b, a)} \int_a^{a+\eta(b, a)} f(x) dx \leq \frac{f(a) + f(b)}{2}.$$

Now, we prove an inequality of Hermite–Hadamard type for h -preinvex fuzzy processes. But first we need the following assumption regarding the function η which is due to S.R. Mohan and S.K. Neogy [1995]:

Condition C. Let $X \subseteq R$ be an open invex subset with respect to η . For any $x, y \in X$ and any $t \in [0, 1]$

$$\begin{aligned} \eta(y, y + t\eta(x, y)) &= -t\eta(x, y) \\ \eta(x, y + t\eta(x, y)) &= (1-t)\eta(x, y). \end{aligned}$$

Note that every $x, y \in X$ and every $t_1, t_2 \in [0, 1]$ from the above Condition C, we have

$$\eta(y + t_2\eta(x, y), y + t_1\eta(x, y)) = (t_2 - t_1)\eta(x, y).$$

Theorem 4.1. Let $F : [a, a + \eta(b, a)] \rightarrow \mathcal{F}_K(R^n)$ be *h*-preinvex and integrably bounded fuzzy process, with $a < a + \eta(b, a)$. Then, if $h\left(\frac{1}{2}\right) \neq 0$ and the Condition C is fulfilled we have

$$\begin{aligned} \{F(a) + F(b)\} \int_0^1 h(t)dt \\ \subseteq \frac{1}{\eta(b, a)} \int_a^{a+\eta(b, a)} F(x)dx \subseteq \frac{1}{2h\left(\frac{1}{2}\right)} F(a + \eta(b, a)). \end{aligned}$$

Proof. Since F is *h*-preinvex fuzzy process we have that

$$h(t)F(a) + h(1 - t)F(b) \subseteq F(a + t\eta(b, a))$$

for all $t \in (0, 1)$. Integrating on $[0, 1]$, we obtain

$$\begin{aligned} \int_0^1 F(a + \eta(b, a))dt \supseteq \int_0^1 h(t)F(a) + h(1 - t)F(b)dt \\ = \{F(a) + F(b)\} \cdot \int_0^1 h(t)dt. \end{aligned}$$

Making the change of variable $x = a + t \eta(b, a)$ we obtain

$$\frac{1}{\eta(b, a)} \int_a^{a+\eta(b, a)} F(x)dx \supseteq \{F(a) + F(b)\} \int_0^1 h(t)dt.$$

To prove the second inclusion, we observe that from the Condition C it follows that

$$\begin{aligned} F\left(a + \frac{1}{2}\eta(b, a)\right) \\ = F\left(a + t\eta(b, a) + \frac{1}{2}\eta(a + (1 - t)\eta(b, a), a + t\eta(b, a))\right) \\ \supseteq h\left(\frac{1}{2}\right) F(a + t\eta(b, a)) + F(a + (1 - t)\eta(b, a)). \end{aligned}$$

Integrating this inclusion on $[0,1]$ we get

$$\begin{aligned} \int_0^1 F\left(a + \frac{1}{2}\eta(b, a)\right) dt \\ \supseteq h\left(\frac{1}{2}\right) \left[\int_0^1 F\left(a + \frac{1}{2}\eta(b, a)\right) dt \right. \\ \left. + \int_0^1 F\left(a + (1-t)\eta(b, a)\right) dt \right]. \end{aligned}$$

Since

$$\begin{aligned} \int_0^1 F\left(a + t\eta(b, a)\right) dt &= \int_0^1 F\left(a + (1-t)\eta(b, a)\right) dt \\ &= \frac{1}{\eta(b, a)} \int_a^{a+\eta(b, a)} F(x) dx, \end{aligned}$$

it follows that

$$\frac{1}{\eta(b, a)} \int_a^{a+\eta(b, a)} F(x) dx \subseteq \frac{1}{2h\left(\frac{1}{2}\right)} F\left(a + \frac{1}{2}\eta(b, a)\right)$$

what completes the proof.

Remark 4.1. If $\eta(b, a) = b - a$ and $h(t) = t^s$ then our inequality reduces to the inequality for s -convex fuzzy processes obtained by R. Osuna-Gomez et al. [2004].

Now, for a mapping $f: [a, b] \rightarrow R$ let us define a mapping

$$H(t) = \frac{1}{b-a} \int_a^b f\left(tx + (1-t)\frac{a+b}{2}\right) dx.$$

Some properties of this mapping for convex function are given in [Dragomir 1992]. We extend these properties for h -convex fuzzy processes. Let $F: [a, b] \rightarrow \mathcal{F}_C(R^n)$ be an integrably bounded fuzzy process and define

$$H_F(t) = \frac{1}{b-a} \int_a^b F\left(tx + (1-t)\frac{a+b}{2}\right) dx$$

for $t \in [0, 1]$.

Theorem 4.2. Let F be a *h*-convex integrably bounded fuzzy process on an interval $[a, b]$. Then H_F is *h*-convex on $[0, 1]$ and

$$H_F(t) \subseteq \frac{1}{2h\left(\frac{1}{2}\right)} H_F(0), \forall t \in [0, 1].$$

Proof. The *h*-convexity of the mapping H_F is a consequence of the *h*-convexity of the mapping F . Namely, for $t_1, t_2 \in [0, 1]$ and $\alpha, \beta \geq 0$ with $\alpha + \beta = 1$ we have:

$$\begin{aligned} & H_F(\alpha t_1 + \beta t_2) \\ &= \frac{1}{b-a} \int_a^b F\left((\alpha t_1 + \beta t_2)x + (1 - (\alpha t_1 + \beta t_2))\frac{a+b}{2}\right) dx \\ &= \frac{1}{b-a} \int_a^b F\left(\alpha(t_1 x + (1-t_1)\frac{a+b}{2}) + \beta(t_2 x + (1-t_2)\frac{a+b}{2})\right) dx \\ &\supseteq \frac{1}{b-a} \int_a^b h(\alpha)F\left(t_1 x + (1-t_1)\frac{a+b}{2}\right) dx \\ &\quad + \frac{1}{b-a} \int_a^b h(\beta)F\left(t_2 x + (1-t_2)\frac{a+b}{2}\right) dx \\ &= h(\alpha)H_F(t_1) + h(\beta)H_F(t_2) \end{aligned}$$

what means that H_F is *h*-convex.

Now, let $t \in [0, 1]$. Taking $y = t x + (1-t)\frac{a+b}{2}$ we obtain

$$H_F = \frac{1}{d-c} \int_c^d F(y) dy$$

where $c = ta + (1-t)\frac{a+b}{2}$ and $d = tb + (1-t)\frac{a+b}{2}$. By Theorem 4.1 for $\eta(d, c) = d - c$ we have

$$\begin{aligned} \frac{1}{d-c} \int_c^d F(y) dy &\subseteq \frac{1}{2h\left(\frac{1}{2}\right)} F\left(\frac{c+d}{2}\right) = \frac{1}{2h\left(\frac{1}{2}\right)} F\left(\frac{a+b}{2}\right) \\ &= \frac{1}{2h\left(\frac{1}{2}\right)} H_F(0). \end{aligned}$$

what completes the proof.

References

- Aumann R.I., 1965, *Integrals of set-valued functions*, Journal of Mathematical Analysis and Applications, vol. 12, pp. 1–12.
- Bector C.R., Singh C., 1991, *B-vex functions*, Journal of Optimization Theory and Applications, vol. 71, pp. 237–253.
- Breckner W.W., 1978, *Stetigkeitsaussagen für eine Klasse verallgemeinerter Konvexer Funktionen in topologischen linearen Räumen*, Publications de l'Institut Mathématique (Beograd) 23, pp. 13–20.
- Breckner W.W., 1993, *Continuity of generalized convex and generalized concave set-valued functions*, Revue d'Analyse Numérique et de Théorie de l'Approximation, vol. 22, pp. 39–51.
- Brunovsky P., 1968, *On the necessity of a certain convexity condition for lower closure of control problems*, SIAM Journal on Control, vol. 6, pp. 174–185.
- Chalco-Cano Y., Rojas-Medar M.A., 2004, *s-Convex fuzzy processes*, Computers & Mathematics with Applications, vol. 47, pp. 1411–1418.
- Coutat P., 1996, *More on continuous closed convex sets*, Optimization, vol. 37, pp. 99–112.
- Debreu G., 1983, *Mathematical Economics: Twenty Papers of Gerard Debreu*, Cambridge University Press, Cambridge–New York.
- Dragomir S.S., 1992, *Two mappings in connection to Hadamard's inequalities*, Journal of Mathematical Analysis and Applications, vol. 167, pp. 49–56.
- Dragomir S.S., Fitzpatrick S., 1999, *The Hadamard's inequality for s-convex functions in the second sense*, Demonstratio Mathematica, vol. 32, no. 4, pp. 687–696.
- Hanson M.A., 1981, *On sufficiency of Kuhn–Tucker Conditions*, Journal of Mathematical Analysis and Applications, vol. 80, pp. 545–550.
- Hermes H., 1968, *Calculus of set valued functions and control*, Journal of Mathematics and Mechanics, vol. 18, pp. 47–60.
- Lowen R., 1980, *Convex fuzzy sets*, Fuzzy Sets and Systems, vol. 3, pp. 291–310.
- Matłoka M., 1999, *Some properties of Φ -convex fuzzy processes*, Journal of Fuzzy Mathematics, vol. 7, pp. 701–706.
- Matłoka M., 2000, *Convex fuzzy processes*, Fuzzy Sets and Systems, vol. 110, pp. 109–114.
- Matłoka M., 2012, *A note on $(\Phi E_1, \Phi E_2)$ -convex fuzzy processes*, Advances in Fuzzy Systems, vol. 2012, article ID 290845.
- Mohan S.R., Neogy S.K., 1995, *On invex sets and preinvex functions*, Journal of Mathematical Analysis and Applications, vol. 189, pp. 901–908.
- Noor M.S., 2009, *Hadamard integral inequalities for product of two preinvex functions*, Nonlinear Analysis Forum, vol. 14, pp. 167–173.
- Osuna-Gómez R., Jimenez-Gamero M.D., Chalco-Cano Y., Rojas-Medar M.A., 2004, *Hadamard and Jensen inequalities for s-convex fuzzy processes*, [in:] Lopez-Diaz M., Gil M.A., Grzegorzewski P., Hryniewicz O., Lawry L. (eds.) *Soft Methodology and Random Information Systems*, vol. 1, Springer, Heidelberg, pp. 645–652.
- Pini R., Singh Ch., 1997, *(Φ_1, Φ_2) -convexity*, Optimization, vol. 40, pp. 103–120.
- Puri M.L., Ralescu D.A., 1987, *Fuzzy random variables*, Journal of Mathematical Analysis and Applications, vol. 114, pp. 301–317.
- Rockafellar R.T., 1970, *Convex Analysis*, Princeton University Press, Princeton, NJ.
- Sarikaya M.Z., Saglam A., Yildirim H., 2008, *On some Hadamard type inequalities for h-convex functions*, Journal of Mathematical Inequalities, vol. 2, pp. 335–341.

- Syau Y.R., Low C.Y., Wu T.H., 2002, *A note on convex fuzzy processes*, Applied Mathematics Letters, vol. 15, pp. 193–196.
- Varošanec S., 2007, *On h -convexity*, Journal of Mathematical Analysis and Applications, vol. 326, pp. 303–311.
- Weir T., Mond B., 1988, *Preinvex functions in multiobjective optimization*, Journal of Mathematical Analysis and Applications, vol. 136, pp. 29–38.
- Youness E.A., 1999, *E -convex sets, E -convex functions and E -convex programming*, Journal of Optimization Theory and Applications, vol. 102, pp. 439–450.