

Propagation properties of anomalous hollow vortex beam in uniaxial crystal orthogonal to the optical axis

DAJUN LIU*, YAPOCHUAN WANG, GUIQIU WANG

Department of Physics, Dalian Maritime University, Dalian, 116026, China

*Corresponding author: liudajun@dlnu.edu.cn

The analytical equation of an anomalous hollow vortex beam propagating in a uniaxial crystal orthogonal to the optical axis is derived, and the intensity and phase properties of this beam propagating in the uniaxial crystal are illustrated using numerical examples. The influences of topological charge M and the ratio of refractive index n_e/n_o of a uniaxial crystal on the normalized intensity distribution and phase distribution for anomalous hollow vortex beam are discussed and analyzed in detail.

Keywords: uniaxial crystal, anomalous hollow vortex beam, laser propagation.

1. Introduction

Recently, the propagation properties of various laser beams in a uniaxial crystal have been widely studied because the uniaxial crystal has the application in the design of wave plates and optical devices [1]. Until now, based on the propagation theory of laser beams in a uniaxial crystal [2–4], the evolution properties of various laser beams propagating in a uniaxial crystal have been widely illustrated, such as those of Laguerre–Gauss and Bessel–Gauss beams [5], dark hollow beam [6], anomalous hollow beam [7], Hermite-cosine-Gaussian beam [8], partially polarized partially coherent beam [9], partially coherent flat-topped beam [10], Lorentz and Lorentz–Gauss beams [11], partially coherent Lorentz beam [12], twisted Gaussian Schell-model beam [13], Airy beam [14], partially coherent anomalous hollow beam [15], elliptical Gaussian vortex beam [16], four-petal Gaussian vortex beam [17], Laguerre–Gaussian correlated Schell-model beam [18], partially coherent four-petal Gaussian vortex beams [19], flat-topped vortex hollow

beam [20], partially coherent flat-topped vortex hollow beam [21], multi-Gaussian Schell model beam [22] and rectangular multi-Gaussian Schell model beam [23].

On the other hand, new laser beams have been introduced to describe the optical field of a laser output. Among the laser beams, a new dark hollow beam called an anomalous hollow beam has been produced [24]. And the properties of the anomalous hollow beam propagating in the uniaxial crystal, turbulent atmosphere, optical system and free space have been widely studied [7, 25–27]. Based on the model of the anomalous hollow beam, the anomalous hollow beam with an orbital angular momentum has also been introduced and studied [28]. Until now, there have been no reports about the anomalous hollow vortex beam propagating in the uniaxial crystal orthogonal to the optical axis. In this paper, our aim is to study the intensity and phase properties of the anomalous hollow vortex beam propagating in the uniaxial crystal orthogonal to the optical axis.

2. Propagation theory

In the Cartesian coordinate system, the z -axis is set to be the propagation axis, and the electric field of an anomalous hollow vortex beam at the source plane $z = 0$ can be expressed as [28]

$$E(\mathbf{r}_0, 0) = \left(-2 + \frac{8x_0^2}{w_{0x}^2} + \frac{8y_0^2}{w_{0y}^2} \right) \exp \left(-\frac{x_0^2}{w_{0x}^2} - \frac{y_0^2}{w_{0y}^2} \right) (x_0 + iy_0)^M \quad (1)$$

where $\mathbf{r}_0 = (x_0, y_0)$ denotes the position vector at the source plane $z = 0$; M is the topological charge; w_{0x} and w_{0y} are the beam waist widths of an astigmatic Gaussian beam in x and y directions, respectively. When $w_{0x} = w_{0y}$, Eq. (1) reduces to a circular anomalous hollow vortex beam.

Recalling the following formula [29]

$$(x + iy)^M = \sum_{l=0}^M \frac{M! i^l}{l!(M-l)!} x^{M-l} y^l \quad (2)$$

the electric field of an anomalous hollow vortex beam can be written as

$$E(\mathbf{r}_0, 0) = \sum_{l=0}^M \frac{M! i^l}{l!(M-l)!} x_0^{M-l} y_0^l \left(-2 + \frac{8x_0^2}{w_{0x}^2} + \frac{8y_0^2}{w_{0y}^2} \right) \exp \left(-\frac{x_0^2}{w_{0x}^2} - \frac{y_0^2}{w_{0y}^2} \right) \quad (3)$$

Under paraxial approximation, we assume that the optical axis of the uniaxial crystal coincides with the x -axis of the Cartesian coordinate system, and the beam propagates along the z -axis, thus the electric field of the beam propagating in the uniaxial crystal orthogonal to the optical axis can be expressed as:

$$E_x(\mathbf{r}, z) = \exp(ikn_e z) \frac{kn_o}{2\pi iz} \times \int dx_0 dy_0 \exp\left\{-\frac{k}{2izn_e} \left[n_o^2(x-x_0)^2 + n_e^2(y-y_0)^2\right]\right\} E_x(\mathbf{r}_0, 0) \quad (4a)$$

$$E_y(\mathbf{r}, z) = \exp(ikn_o z) \frac{kn_o}{2\pi iz} \times \int dx_0 dy_0 \exp\left\{-\frac{kn_o}{2iz} \left[(x-x_0)^2 + (y-y_0)^2\right]\right\} E_y(\mathbf{r}_0, 0) \quad (4b)$$

where $k = 2\pi/\lambda$ is the wavenumber with λ denoting the wavelength; $\mathbf{r} = (x, y)$ is the position vectors at the observation plane. From Eq. (4a), it can be seen that the only x component of the beam propagating in uniaxial crystals undergoes diffraction spreading asymmetry in the x - y plane due to the influences of the uniaxial crystal. Therefore, only the x -polarized anomalous hollow vortex beam propagating in uniaxial crystals orthogonal to the optical axis is analyzed in this work.

Substituting Eq. (3) into Eq. (4a), and recalling the following formulae [29]:

$$\int_{-\infty}^{+\infty} x^n \exp(-px^2 + 2qx) dx = n! \exp\left(\frac{q^2}{p}\right) \left(\frac{q}{p}\right)^n \sqrt{\frac{\pi}{p}} \sum_{k=0}^{[n/2]} \frac{1}{k!(n-2k)!} \left(\frac{p}{4q^2}\right)^k \quad (5)$$

$$H_n(x) = \sum_{k=0}^{[n/2]} \frac{(-1)^k n!}{k!(n-2k)!} (2x)^{n-2k} \quad (6)$$

The analytical propagation equation of the anomalous hollow vortex beam propagating in the uniaxial crystal orthogonal to the optical axis can be derived as

$$E(x, y, z) = \frac{kn_o}{2\pi iz} \exp(ikn_e z) \exp\left(-\frac{kn_o^2}{2izn_e} x^2 - \frac{kn_e}{2iz} y^2\right) \times \sum_{l=0}^M \frac{M! i^l}{l!(M-l)!} \left(-2I_1 + \frac{8}{w_{0x}^2} I_2 + \frac{8}{w_{0y}^2} I_3\right) \quad (7)$$

where

$$I_1 = 2^{-M} i^M \sqrt{\frac{\pi}{a_x}} \sqrt{\frac{\pi}{a_y}} \left(\frac{1}{\sqrt{a_x}}\right)^{M-l} \left(\frac{1}{\sqrt{a_y}}\right)^l \exp\left(\frac{c_x^2}{a_x}\right) \exp\left(\frac{c_y^2}{a_y}\right) \times H_{M-l}\left(-\frac{ic_x}{\sqrt{a_x}}\right) H_l\left(-\frac{ic_y}{\sqrt{a_y}}\right) \quad (8)$$

$$\begin{aligned}
I_2 = & 2^{-M-2} i^{M+2} \sqrt{\frac{\pi}{a_x}} \sqrt{\frac{\pi}{a_y}} \left(\frac{1}{\sqrt{a_x}} \right)^{M-l+2} \left(\frac{1}{\sqrt{a_y}} \right)^l \exp\left(\frac{c_x^2}{a_x}\right) \exp\left(\frac{c_y^2}{a_y}\right) \\
& \times H_{M-l+2}\left(-\frac{ic_x}{\sqrt{a_x}}\right) H_l\left(-\frac{ic_y}{\sqrt{a_y}}\right)
\end{aligned} \tag{9}$$

$$\begin{aligned}
I_3 = & 2^{-M-2} i^{M+2} \sqrt{\frac{\pi}{a_x}} \sqrt{\frac{\pi}{a_y}} \left(\frac{1}{\sqrt{a_x}} \right)^{M-l} \left(\frac{1}{\sqrt{a_y}} \right)^{l+2} \exp\left(\frac{c_x^2}{a_x}\right) \exp\left(\frac{c_y^2}{a_y}\right) \\
& \times H_{M-l}\left(-\frac{ic_x}{\sqrt{a_x}}\right) H_{l+2}\left(-\frac{ic_y}{\sqrt{a_y}}\right)
\end{aligned} \tag{10}$$

with

$$a_x = \frac{1}{w_{0x}^2} + \frac{kn_0^2}{2izn_e} \tag{11a}$$

$$c_x = \frac{kn_0^2}{2izn_e} x \tag{11b}$$

$$a_y = \frac{1}{w_{0y}^2} + \frac{kn_e}{2iz} \tag{12a}$$

$$c_y = \frac{kn_e}{2iz} y \tag{12b}$$

The intensity and phase distributions of the anomalous hollow vortex beam propagating in the uniaxial crystal orthogonal to the optical axis can be illustrated and analyzed using the derived Eqs. (7)–(12).

3. Numerical examples and analysis

In this section, the intensity and phase properties of the anomalous hollow vortex beam propagating in the uniaxial crystal orthogonal to the optical axis are illustrated and analyzed using numerical examples. Unless specified otherwise in captions of figures, the parameters of the uniaxial crystal and the anomalous hollow vortex beam are chosen as $n_0 = 2.616$, $\lambda = 1.064 \mu\text{m}$, $w_{0x} = 10 \mu\text{m}$, $M = 2$ and Rayleigh length $z_R = \pi w_{0x}^2 n_0 / \lambda$.

The contour graphs of the normalized intensity of the anomalous hollow vortex beam propagating in the uniaxial crystal orthogonal to the optical axis for different n_e/n_0 are illustrated in Figs. 1 and 2. It can be found that the anomalous hollow vortex beam keeps its initial central dark centre as the propagation distance increases, and the width




Fig. 1. The contour graphs of the normalized intensity of anomalous hollow vortex beam in uniaxial crystal with $n_e/n_o = 1.1$, $w_{0y} = 10 \mu\text{m}$, and $z = z_R$ (a), $z = 2z_R$ (b), $z = 5z_R$ (c), $z = 10z_R$ (d).




Fig. 2. The contour graphs of the normalized intensity of anomalous hollow vortex beam in uniaxial crystal with $n_e/n_o = 1.3$, $w_{0y} = 10 \mu\text{m}$, and $z = z_R$ (a), $z = 2z_R$ (b), $z = 5z_R$ (c), $z = 10z_R$ (d).

of beam will spread rapidly as the propagation distance increases. The beam propagating in the uniaxial crystal with smaller n_e/n_o (Fig. 1) can almost keep its circular symmetry, and the beam propagating in the uniaxial crystal with larger n_e/n_o (Fig. 2) will evolve into an elliptical symmetry beam with a central dark centre. In Fig. 3, the propagation of the anomalous hollow vortex beam with elliptical symmetry in the uni-




Fig. 3. The contour graphs of the normalized intensity of anomalous hollow vortex beam in uniaxial crystal with $n_e/n_o = 1.1$, $w_{0y} = 13 \mu\text{m}$, and $z = z_R$ (a), $z = 2z_R$ (b), $z = 5z_R$ (c), $z = 10z_R$ (d).




Fig. 4. Cross-section of anomalous hollow vortex beam in uniaxial crystal for different n_e/n_o at the propagation distance $z = 5z_R$ along x axis (a), and along y axis (b).




Fig. 5. Cross-section ($y = 0$) of anomalous hollow vortex beam in uniaxial crystal with $n_e/n_o = 1.1$ for different M ; $z = z_R$ (a), and $z = 5z_R$ (b).

axial crystal with $n_e/n_o = 1.1$ is given. It can be found that the anomalous hollow vortex beam with elliptical symmetry propagating in the uniaxial crystal with smaller n_e/n_o will keep its initial spot pattern and spread as the propagation distance increases.

Figures 4 and 5 show the cross-section of the anomalous hollow vortex beam propagating in the uniaxial crystal for different n_e/n_o and M , respectively. As can be seen,




Fig. 6. The phase of anomalous hollow vortex beam propagating in uniaxial crystal with $n_e/n_o = 1.1$ at the propagation distance $z = z_R$ for $M = 1$ (a), $M = 2$ (b), $M = 3$ (c), and $M = 4$ (d).




Fig. 7. The phase of anomalous hollow vortex beam propagating in uniaxial crystal with $n_e/n_o = 1.1$; $z = z_R$ (a), and $z = 5z_R$ (b).




Fig. 8. The phase of anomalous hollow vortex beam propagating in uniaxial crystal with $n_e/n_o = 1.3$; $z = z_R$ (a), and $z = 5z_R$ (b).

the beam propagating in the uniaxial crystal will spread faster along the x axis as n_e/n_o increases, and spread slower along the y axis as n_e/n_o increases. The reason for this is that the beam propagation in the uniaxial crystal along the x axis and the y axis has different Rayleigh length, and the Rayleigh length along the x axis $\pi w_{0x}^2 n_o / \lambda$ is smaller than the Rayleigh length along the y axis $\pi w_{0y}^2 n_e / \lambda$ as n_e/n_o increases. And the beam with larger M will have a larger central dark centre.

The phases of the anomalous hollow vortex beam propagating in the uniaxial crystal for different M and n_e/n_o are given in Figs. 6–8. It is found that the beam with different topological charge M will keep its number of coherent vortex unchanged, while the partially coherent vortex beam will have the changed number of coherent vortices [21] as the propagation distance increases. When $M=2$, the phase distributions of the anomalous hollow vortex beam propagating in the uniaxial crystal with smaller n_e/n_o have the double clock wise spiral distribution (Fig. 7); and as n_e/n_o of the uniaxial crystal increases, the phase distributions become irregular with the propagation distance z increasing, and the phase distributions become more irregular in the x direction than the y direction due to the $n_e/n_o > 1$ of the uniaxial crystal, which is caused by the different Rayleigh lengths along the x axis and y axis.

4. Conclusions

In this paper, the analytical equation of an anomalous hollow vortex beam propagating in the uniaxial crystal orthogonal to the optical axis is derived, and the propagation properties of the beam are analyzed using numerical examples. The results show that the beam propagation in the uniaxial crystal will keep its initial central dark centre and spread as the propagation distance increases. The beam propagating in the uniaxial crystal will spread faster along the x axis as n_e/n_o increases, and spread slower along the y axis as n_e/n_o increases. The phase distributions of the beam propagating in the uniaxial crystal with larger n_e/n_o are more irregular in the x direction than y direction due to the difference in Rayleigh lengths along the x direction and the y direction.

Acknowledgements – This work was supported by National Natural Science Foundation of China (11604038, 11404048, 11375034), Natural Science Foundation of Liaoning Province (201602062, 201602061) and the Fundamental Research Funds for the Central Universities (3132018235, 3132018236).

References

- [1] WOLF M.B.E., *Principle of Optics*, Pergamon Press, 1999.
- [2] CIATTONI A., CROSIGNANI B., DI PORTO P., *Vectorial theory of propagation in uniaxially anisotropic media*, [Journal of the Optical Society of America A 18\(7\), 2001, pp. 1656–1661.](#)
- [3] CIATTONI A., CROSIGNANI B., DI PORTO P., *Vectorial analytical description of propagation of a highly nonparaxial beam*, [Optics Communications 202\(1–3\), 2002, pp. 17–20.](#)
- [4] CIATTONI A., CINCOTTI G., PROVENZIANI D., PALMA C., *Paraxial propagation along the optical axis of a uniaxial medium*, [Physical Review E 66\(3\), 2002, article ID 036614.](#)
- [5] CINCOTTI G., CIATTONI A., PALMA C., *Laguerre–Gauss and Bessel–Gauss beams in uniaxial crystals*, [Journal of the Optical Society of America A 19\(8\), 2002, pp. 1680–1688.](#)
- [6] DAJUN LIU, ZHONGXIANG ZHOU, *Various dark hollow beams propagating in uniaxial crystals orthogonal to the optical axis*, [Journal of Optics A: Pure and Applied Optics 10\(9\), 2008, article ID 095005.](#)
- [7] DAJUN LIU, ZHONGXIANG ZHOU, *Propagation properties of anomalous hollow beam in uniaxial crystals orthogonal to the optical axis*, [Optics and Laser Technology 41\(7\), 2009, pp. 877–884.](#)
- [8] BIN TANG, *Hermite-cosine-Gaussian beams propagating in uniaxial crystals orthogonal to the optical axis*, [Journal of the Optical Society of America A 26\(12\), 2009, pp. 2480–2487.](#)
- [9] LIU D.J., ZHOU Z.X., *Propagation of partially polarized, partially coherent beams in uniaxial crystals orthogonal to the optical axis*, [The European Physical Journal D 54\(1\), 2009, pp. 95–101.](#)
- [10] DAJUN LIU, ZHONGXIANG ZHOU, *Propagation of partially coherent flat-topped beams in uniaxial crystals orthogonal to the optical axis*, [Journal of the Optical Society of America A 26\(4\), 2009, pp. 924–930.](#)
- [11] CHENGLIANG ZHAO, YANGJIAN CAI, *Paraxial propagation of Lorentz and Lorentz–Gauss beams in uniaxial crystals orthogonal to the optical axis*, [Journal of Modern Optics 57\(5\), 2010, pp. 375–384.](#)
- [12] DAJUN LIU, HONGMING YIN, GUIQIU WANG, YAOUCHUAN WANG, *Propagation properties of a partially coherent Lorentz beam in uniaxial crystal orthogonal to the optical axis*, [Journal of the Optical Society of America A 34\(6\), 2017, pp. 953–960.](#)
- [13] LINA ZHANG, YANGJIAN CAI, *Evolution properties of a twisted Gaussian Schell-model beam in a uniaxial crystal*, [Journal of Modern Optics 58\(14\), 2011, pp. 1224–1232.](#)
- [14] GUOQUAN ZHOU, RUIPIN CHEN, XIUXIANG CHU, *Propagation of Airy beams in uniaxial crystals orthogonal to the optical axis*, [Optics Express 20\(3\), 2012, pp. 2196–2205.](#)

- [15] XIN WANG, XINGYUAN LU, CHENCHEN ZHAO, KUILONG WANG, CHENGLIANG ZHAO, YANGJIAN CAI, *Propagation properties of partially coherent anomalous hollow beams in uniaxial crystals*, [Journal of Modern Optics 61\(8\), 2014, pp. 688–696.](#)
- [16] XUN WANG, ZHIRONG LIU, DAOMU ZHAO, *Nonparaxial propagation of elliptical Gaussian vortex beams in uniaxial crystal orthogonal to the optical axis*, [Journal of the Optical Society of America A 31\(10\), 2014, pp. 2268–2274.](#)
- [17] DAJUN LIU, HE WANG, YAOCHUAN WANG, HONGMING YIN, *Evolution properties of four-petal Gaussian vortex beam propagating in uniaxial crystals orthogonal to the optical axis*, [The European Physical Journal D 69\(9\), 2015, article ID 218.](#)
- [18] ZIREN ZHU, LIN LIU, FEI WANG, YANGJIAN CAI, *Evolution properties of a Laguerre–Gaussian correlated Schell-model beam propagating in uniaxial crystals orthogonal to the optical axis*, [Journal of the Optical Society of America A 32\(3\), 2015, pp. 374–380.](#)
- [19] DAJUN LIU, YAOCHUAN WANG, HONGMING YIN, *Evolution properties of partially coherent four-petal Gaussian vortex beams propagating in uniaxial crystals orthogonal to the optical axis*, [Journal of the Optical Society of America A 32\(9\), 2015, pp. 1683–1690.](#)
- [20] DAJUN LIU, YAOCHUAN WANG, GUIQIU WANG, HONGMING YIN, *Propagation properties of flat-topped vortex hollow beam in uniaxial crystals orthogonal to the optical axis*, [Optik – International Journal for Light and Electron Optics 127\(19\), 2016, pp. 7842–7851.](#)
- [21] DAJUN LIU, GUIQIU WANG, XIXIAN LUO, HONGMING YIN, YAOCHUAN WANG, *Evolution properties of a partially coherent flat-topped vortex hollow beam propagating in uniaxial crystals orthogonal to the optical axis*, [Journal of the Optical Society of Korea 20\(6\), 2017, pp. 686–693.](#)
- [22] XINGYUAN LU, YAN SHEN, XINLEI ZHU, CHENGLIANG ZHAO, YANGJIAN CAI, *Evolution properties of multi-Gaussian Schell model beams propagating in uniaxial crystal orthogonal to the optical axis*, [Optica Applicata 46\(1\), 2016, pp. 19–34.](#)
- [23] YONGHUA MAO, ZHANGRONG MEI, *Propagation properties of the rectangular multi-Gaussian Schell-model beams in uniaxial crystals orthogonal to the optical axis*, [IEEE Photonics Journal 9\(2\), 2017, article ID 6100410.](#)
- [24] YANGJIAN CAI, ZHAOYING WANG, QIANG LIN, *An alternative theoretical model for an anomalous hollow beam*, [Optics Express 16\(19\), 2008, pp. 15254–15267.](#)
- [25] YANGJIAN CAI, EYYUBOĞLU H.T., BAYKAL Y., *Propagation properties of anomalous hollow beams in a turbulent atmosphere*, [Optics Communications 281\(21\), 2008, pp. 5291–5297.](#)
- [26] KUILONG WANG, CHENGLIANG ZHAO, BIJUN XU, *Propagation of anomalous hollow beam through a misaligned first-order optical system*, [Optics and Laser Technology 42\(8\), 2010, pp. 1218–1222.](#)
- [27] KUILONG WANG, CHENGLIANG ZHAO, *Nonparaxial propagation of a vectorial apertured off-axis anomalous hollow beam*, [Optics Communications 334, 2015, pp. 280–286.](#)
- [28] CHENCHEN ZHAO, XIN WANG, CHENGLIANG ZHAO, KUILONG WANG, YANGJIAN CAI, *Statistical properties of an anomalous hollow beam with orbital angular momentum*, [Journal of Modern Optics 62\(3\), 2015, pp. 179–185.](#)
- [29] JEFFREY A., HUI-HUI DAI, *Handbook of Mathematical Formulas and Integrals*, 4th Ed., Academic Press, 2008.

*Received May 23, 2017
in revised form June 29, 2017*